

Features Influencing Older Adults' Quality of Life and Their Association With Loneliness and Activities of Daily Living: A Cross-sectional Study in Iran

Zahra Rostami¹, Mehdi Abbasian^{2,3}, Ehsan Sarbazi^{4*}, Hassan Soleimanpour⁵, Hadi Mostafaei⁶, Masouma Ghasemi², Mohammad Sadra Nemati⁷, Hosein Azizi⁸

¹Student Research Committee, Zanjan University of Medical Sciences, Zanjan, Iran

²Student Research Committee, Tabriz University of Medical Sciences, Tabriz, Iran

³Research Center for Integrative Medicine in Aging, Aging Research Institute, Tabriz University of Medical Sciences, Tabriz, Iran

⁴Road Traffic Injury Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

⁵Emergency and Trauma Care Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

⁶Resident of Urology, Medical University of Vienna and Research Center for EBM, Vienna, Austria

⁷Young Researcher and Elite Club, Tehran Medical Sciences, Islamic Azad University, Tehran, Iran

⁸Women's Reproductive Health Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

Article History:

Received: May 31, 2023

Accepted: December 10, 2023

ePublished: December 20, 2023

*Corresponding Author:

Ehsan Sarbazi,

Email: Ehsansarbazi20@gmail.com

Abstract

Objectives: This study was conducted to assess the features that affect the quality of life (QoL) of older adults and how they are linked to their ability to perform activities of daily living (ADL) and feelings of loneliness.

Design: A random sampling method was employed for this cross-sectional study.

Setting (s): Tabriz, East Azerbaijan, Iran.

Participants: Home-dwelling older adults.

Outcome measures: Data collection tools included the SF-12 QoL, ADL, Loneliness Scale, and demographic information. Relationships between loneliness, ADL, and QoL were examined using a multiple linear regression model by means of the backward method.

Results: On average, the age of the participants was 67.71 years with a standard deviation (SD) of 7.13 years. Of all these older adults, 209 (67%) were aged 60-69 years, 270 (86%) were married, and 155 (49.4%) had elementary education. Mean \pm SD for the physical and mental components of QoL were obtained at 49.35 ± 9.45 and 48.38 ± 11.65 , respectively. Emotional loneliness had a significant reversed association ($\beta = -0.38$, $P < 0.01$) with Mental Component Summary (MCS), while social loneliness had a significant positive association ($\beta = 0.27$, $P = 0.03$) with MCS. Furthermore, dependent ADL had a negative correlation ($\beta = -0.23$; $P < 0.01$) with Physical Component Summary (PCS). Moreover, age, gender, and educational status had a statistically significant relationship with QoL.

Conclusions: Given the significant negative impact of loneliness and low ADL scores, it is imperative to develop efficient interventions and modify existing procedures to improve the QoL for older people. Individual-level interventions and societal-level interventions need to be developed and applied among this population. Additionally, national health policymakers may prioritize loneliness as a significant concern for public health.

Keywords: Activities of daily living, Loneliness, Quality of life, Behavioral epidemic, Epidemiology

Introduction

The critical concern of quality of life (QoL) has been brought to attention due to the rapid rise in life expectancy and the aging population.¹ According to the World Health Organization (WHO), an individual's evaluation of their own life is known as QoL, which is influenced by their cultural background, values, objectives, expectations, benchmarks, and worries.² Based on a systematic assessment of the aspect of QoL in older adults³ and considering our country's conditions such as extended

family and activity-based work in the last decades, it seems that activities of daily living (ADL) and loneliness have had important roles in the QoL of older adults.

The results of a review study showed that 42% of the Iranian older people have low, 50% have moderate, and 7% of them have high QoL.⁴ Receiving education on a healthy lifestyle, which includes engaging in regular physical activity, maintaining a nutritious diet, and addressing emotional and other health concerns, has a beneficial impact on one's QoL.⁵ Maintaining and enhancing the



QoL for older adults involves prioritizing physical activity and mental health.⁶

ADL is a series of basic activities performed to maintain and gain independence in daily life such as getting dressed, eating, and doing personal hygiene.⁷ Functional independence and self-reliance in older adults are significantly influenced by ADL.^{8,9} Further, there seems to be a significant association between older adults' ADL and QoL.¹⁰⁻¹³ Accordingly, limitations and dependency on older people in performing ADL can reduce the QoL and increase the costs for older adults.¹⁴

Investigations on older adults documented one of the essential problems that are less frequently addressed, namely, the experienced loneliness in various societies.^{15,16} They are at risk of loneliness or perceived social isolation (as a modern behavioral epidemic and public health issue) due to decreased interactions, limited capacity, performance, changes in physical health, as well as the death of relatives and friends.¹⁷ Numerous older people describe later-life ages as an isolated period and regard it as an undesirable event.^{18,19}

The results of Berg-Weger and Morley's study demonstrated that a significant number of older adults, around one-third or more, are impacted by loneliness.²⁰ In addition, a study in the United States showed that 17%-57% of older adults have loneliness, in particular individuals who are experiencing anxiety, depression, and dementia.²¹ Furthermore, it has been noted that there exists a notable inverse relationship between ADL and psychological aspects, including feelings of isolation.²²⁻²⁴ There is a different rating of loneliness described as a painful feeling among older adults.²⁵ Berg-Weger and Morley indicated that loneliness as a health issue is not adequately addressed and needs further attention from professionals and social workers.²⁰ Generally speaking, lifestyle modification can greatly affect the capability and independence of aged persons, assisting individuals in managing and regulating the limitations and impairments of bodily systems. Additionally, this practice decreases the likelihood of developing illnesses and disorders later in life.²⁶

Increasing the number of older population and decreasing their accessible resources probably will exacerbate their social support, rehabilitation, and healthcare. These people need to spend the remaining years of their lives in peace while maintaining good physical and mental well-being as well. Additionally, living a long life should be accompanied by a high QoL. Therefore, this research sought to explore the relationship between older adults' QoL and its association with ADL and loneliness.

Methods

Design of the Study and Data Collection

We performed an analytical cross-sectional investigation in Tabriz, East Azerbaijan, Iran, from September to November 2022.

Inclusion and Exclusion Criteria

Older adults (aged 60 and over) who had at least a six-month history of living in Tabriz, were active members of the Comprehensive Health Centers (CHCs), and were accompanied by receiving informed consent met the inclusion criteria. After explaining the design and purpose of the study and obtaining informed consent, the questionnaires were completed by conducting an interview. The interview was done by two trained master students. The exclusion criteria included those who had cognitive and psychological problems such as major depression, Alzheimer's (or other Dementia), bipolar disorders, and schizophrenia. Exclusion criteria were also assessed based on the medical records of CHC.

Sampling

Considering the standard deviation (SD) of the QoL score in the older adult population as 14.4²⁷ and the margin of error as 0.05 at a 95% confidence interval, the sample size was established as nearly 314.

$$n = \frac{Z_{1-\frac{\alpha}{2}}^2 S^2}{d^2}$$

Participants were recruited by randomly selecting a sample from 16 CHC, and about 20 samples were chosen from each CHC with an equal gender ratio.

Questionnaires

Three valid instruments, namely, the Iranian version of the 12-item short form (SF-12) Health Survey, the De Jong Gierveld loneliness scale, the Katz ADL scale, and a demographic checklist were used for data collection.

Quality of Life

The SF-12 Health Survey has two summary measures, including six items in the Physical Component Summary (PCS) and six items in the Mental Component Summary (MCS).²⁸ This questionnaire was validated by Zare et al among older people within two dimensions: the PCS Cronbach's alpha (α)=0.68 and the MCS α =0.71.²⁹ The scores range from 0 to 100. A score of 50 or less on PCS-12 is the cut-off for physical condition. Likewise, a score of 42 or less on MCS-12 may suggest clinical depression.³⁰

Loneliness

The 6-item De Jong Gierveld loneliness scale containing six items has been validated for Iranian older adults, which includes three items dealing with emotional loneliness and three related to social loneliness. This questionnaire has 3-point Likert answers including no, more or less, and yes. Items 2, 3, and 4 have negative and neutral answers and are related to social loneliness, while items 1, 5, and 6 have positive and neutral answers related to emotional loneliness.³¹

Activities of Daily Living

The third scale was the 6-item Katz ADL scale. A previous

study confirmed the validity and reliability of the 6-item ADL scale on Iranian older people with an internal consistency α of 86%.³² This scale includes bathing, getting dressed, transfer, toileting, feeding, and continence. To score this questionnaire, if participants are able to perform an activity, they will get one point and otherwise get zero. The total score ranges between 0 (lack of performance) and 6 (maximum performance), 2 or less indicates severe functional impairment, and a score of 4 shows moderate impairment.³²

Data Analysis

We analyzed the gathered data using descriptive statistics, including frequency, mean, and SD to evaluate the attributes of the participants. The normality of the data was checked using the Kolmogorov-Smirnov test. Moreover, the independent *t* test was used to compare the items of QoL with the characteristics of gender. Additionally, the relationship between the PCS and MCS with the characteristics of older individuals was assessed using Spearman and Pearson's tests. Furthermore, the multiple linear regression model was applied to predict the value of an MCS and PCS based on the value of marital status, age, education, loneliness, and ADL variables using the backward method. This technique can help reduce the chances of overfitting the data, find the best subset of features from a given set of features, and make the linear regression model more interpretable. All statistical examinations were double-sided, and it was considered statistically significant when the *P* value was under 0.05. The statistical analyses were carried out using the Statistical Package for Social Sciences (SPSS 24.0, SPSS Inc., Chicago, USA).

Results

Baseline Demographic Profile of the Participants

A total number of 314 older adults (response rate was about 98.7%), including 163 (51.9%) women and 151 (48.1%) men, participated in this research. This study involved participants with a mean age of 67.71 ± 7.13 years. Of all these older adults, 209 (67%) were aged between 60-69, 270 (86%) were married, and 155 (49.4%) had elementary education (Table 1).

Quality of Life Using SF-12

Table 2 presents the scores of QoL, ADL, and loneliness across gender. The mean \pm SD of PCS and MCS of QoL were 49.35 ± 9.45 and 48.38 ± 11.65 , respectively. Moreover, no significant difference was observed between genders. Furthermore, the comparison of QoL items demonstrated that the physical role ($P=0.02$) and social functioning ($P=0.04$) were significantly higher in men, but the mean emotional loneliness of women was significantly higher ($P=0.01$).

Correlation between Quality of Life, Activities of Daily Living, and Loneliness

The PCS exhibited a positive relationship with marital

status, education, and ADL. To be more specific, the correlation coefficients were 0.20, 0.15, and 0.23, respectively. On the other hand, the PCS showed a negative correlation with age, social loneliness, and emotional loneliness (-0.19, -0.24, and -0.28, respectively). Additionally, the MCS had a negative correlation with age and emotional loneliness (-0.11 and -0.17, respectively), as depicted in Table 3.

Multiple Analysis

Multiple regression analysis demonstrated that emotional loneliness ($\beta = -0.38$, $P < 0.001$) has an inverse significant association, and social loneliness ($\beta = 0.27$, $P = 0.03$) has a significant positive association with MCS, with adjusting the mentioned covariates (Table 4, Model A). Moreover, age had an effective significant association with MCS, that is, age > 80 years illustrated a lower MCS compared to the 60-69 age group ($\beta = -0.26$, $P < 0.01$). In addition, the age group (70-79) showed a lower MCS compared

Table 1. Characteristics of the Study Population (N=314)

Variables	Male No. (%)	Female No. (%)	Total No. (%)	
Age groups	60-69	104 (68.9)	105 (64.4)	209 (66.9)
	70-79	35 (23.2)	44 (27.0)	79 (33.1)
	80+	12 (7.9)	14 (8.6)	26 (8.3)
Marital status	Married	136 (90.1)	134 (82.2)	270 (86.0)
	Widowed	15 (9.9)	29 (17.8)	44 (14.0)
Education	Illiterate	43 (28.5)	62 (38.0)	105 (33.4)
	Elementary	75 (49.7)	80 (49.1)	155 (49.4)
	Senior high school or above	33 (21.9)	21 (12.9)	54 (17.2)

Table 2. Quality of Life, Loneliness, and Activities of Daily Living of Older Participants by Gender

Variable	Male Mean (SD)	Female Mean (SD)	Total Mean (SD)	P Value
PCS-12	49.95(9.0)	48.80(9.8)	49.35(9.4)	0.28
MCS-12	48.93(10.8)	47.93(12.3)	48.35(11.6)	0.21
Concepts of SF-12				
Physical functioning	80.79(28.9)	78.52(30.6)	79.52(29.8)	0.54
Physical role	91.05(27.1)	83.74(34.1)	87.30(31.1)	0.02
Bodily pain	78.64(23.8)	77.30(24.0)	77.77(24.0)	0.56
General health	60.26(22.0)	60.27(24.1)	60.24(23.1)	0.93
Vitality	63.04(23.8)	60.12(26.7)	61.46(25.3)	0.22
Social functioning	73.67(21.7)	68.55(23.8)	71.03(22.9)	0.04
Emotional role	92.71(24.7)	89.57(29.6)	90.79(27.8)	0.30
Mental health	63.18(26.4)	62.69(26.5)	62.88(26.4)	0.83
Independent ADL	142(94.0)	148(90.8)	290(92.4)	0.28
Dependent ADL	9(6.0)	15(9.2)	24(7.6)	
Emotional loneliness	0.41(0.9)	0.72(1.1)	0.58(1.0)	0.01
Social loneliness	0.28(0.7)	0.47(0.9)	0.38(0.8)	0.05

Note. PCS: Physical component summary; MCS: Mental component summary; SD: Standard deviation; ADL: Activities of daily living; Physical role: Role limitations due to physical health; *P* values are based on independent samples *t*-test.

Table 3. The Correlation between the Quality of Life, Activities of Daily Living, and Feelings of Loneliness in Older Adults

	Gender	Age	Marital Status	Education	ADL	Loneliness	
						Social Loneliness	Emotional Loneliness
MCS	0.07	-0.11*	-0.11	0.00	0.09	-0.11	-0.17**
PCS	0.05	-0.19**	0.20**	0.15**	0.23**	-0.24**	-0.28**

Note. ADL: Activities of daily living; MCS: Mental component summary; PCS: Physical component summary; * $P < 0.05$; ** $P < 0.01$.

Table 4. A: Multiple Regression Analysis for Predicting Mental Component Summary of Quality of Life, B: Multiple Regression Analysis for Predicting Physical Component Summary of Quality of Life

		Non-standardized Coefficient B	β Coefficient	P Value
Model A				
Marital (Ref=Married)	Single	7.00	0.21	<0.01
Age (Ref=60-69)	70-79	-3.33	-0.13	0.03
	80+	-10.79	-0.26	<0.01
Education (Ref=Illiterate)	Elementary	-2.93	-0.13	0.05
	\geq High school	0.71	0.02	0.72
Loneliness	Emotional	-4.07	-0.38	<0.01
	Social	3.57	0.27	0.03
Model B				
Marital (Ref=Married)	Single	-4.71	-0.17	<0.01
Age (Ref=60-69)	70-79	-3.03	-0.14	0.01
	80+	-2.22	-0.06	0.26
Education (Ref=Illiterate)	elementary	1.25	0.07	0.25
	\geq High school	3.77	0.15	0.01
ADL (Ref=Independent)	Dependent	-8.16	-0.23	<0.01

Note. (A) Dependent variable=MCS: Mental Component Summary, adjusted $R^2 = 0.113$; (B) Dependent variable=PCS: Physical component summary, adjusted $R^2 = 0.171$; Ref: Reference.

to the reference group ($\beta = -0.13$, $P = 0.03$), and single people had a significant positive association with MCS ($\beta = 0.21$, $P < 0.01$) (A). In the second model (B), the factor that influenced PCS was ADL, which exhibited a significant negative relationship between dependency in ADL and PCS. The beta coefficient was -0.23 , with a P value of less than 0.01. Furthermore, being educated was found to have a higher impact on the PCS compared to illiterates ($\beta = 0.15$, $P = 0.01$). Single people indicated a significant inverse association with PCS compared to their married participants ($\beta = -0.17$, $P < 0.01$). Additionally, the age group (70-79) showed a lower MCS as opposed to the reference group ($\beta = -0.14$, $P = 0.01$), as illustrated in Table 4 (Model A).

Discussion

This study examined the relationship between the loneliness status, ADL, and QoL of older adults in Tabriz. The results demonstrated an association between age, gender, education status, ADL, and loneliness status with QoL. Emotional loneliness had an inverse significant association, and social loneliness had a significant positive association with MCS. Moreover, there was a significant inverse association between dependency in ADL and PCS. Furthermore, being educated was found to have a higher impact on the PCS compared to being an illiterate one. While physical role and social functioning were

significantly higher in men, emotional loneliness was significantly higher in women.

Almost one-third of older people often suffer from loneliness.^{33,34} Loneliness has a significant impact on one's QoL and is linked to disease, widowhood, social isolation, and/or reduced family support.^{35,36} The present study agrees with previous findings and offers additional evidence. Older individuals are more likely to live alone compared to other age groups, and some of them may be less engaged in society, leading to potential negative impacts on their mental well-being and increased social isolation.

The mental performance was greatly affected by the coronavirus disease 2019 (COVID-19) pandemic, especially that of seniors.³⁷⁻⁴⁰ To reduce social isolation, using virtual programs about social services and other online programs was reported to increase the QoL of older adults.⁴¹⁻⁴³ A decrease in family communication and the fear of disease spread caused a rise in emotional loneliness.⁴⁴⁻⁴⁶

Age is another factor that has an influence on QoL among older people.⁴⁷ The results indicated that higher age has an inverse significant association with QoL. This finding agrees with Jivraj and colleagues' findings which showed that QoL tends to weaken as they age, partly because of having poorer health.⁴⁸

Compared to married older adults, single ones had a

higher score on MCS; nonetheless, they had a lower score on PCS in terms of QoL. Married individuals experienced increased anxiety about the possibility of losing their partners during the COVID-19 pandemic, leading to a decrease in their QoL.⁴⁹ Along with this, the prevalence of depression among married people was greater during this period.⁵⁰ Poorolajal reported that the prevalence of aggression among the Iranian adult general population is higher in singles.⁵¹ Furthermore, single older adults had a higher QoL due to having better communication with their network of friends.⁴

ADL was established to be an influential factor in PCS. Therefore, ADL-dependent older adults had eight times lesser scores in the physical field of QoL compared to independent ones. Dependency in ADL reduces the QoL in older people.^{52,53} Older individuals who are capable of independently performing Instrumental Activities of Daily Living (IADL) experience a significantly higher QoL.⁵⁴ The COVID-19 pandemic required older adults to stay at home and limit their social activities, causing the closure of their favorite places such as parks, restaurants, and sports clubs. Consequently, older adults experienced a decrease in their opportunities for social interaction and participation in daily activities, which negatively impacted their PCS of QoL.⁵⁵⁻⁵⁷ Unlike younger adults, physical activity is a daily requirement for older individuals and has a significant impact on different aspects of their health, particularly their physical well-being. Older individuals can enhance their health problems by engaging in physical activities, particularly when they combine aerobic and strength exercises.

However, these findings are limited owing to the use of a cross-sectional design. Due to the nature of the study, it is not possible to examine the time precedence in the relationships between the investigated variables. Moreover, the data of this study were collected during the COVID-19 pandemic, and its data may be different from the non-pandemic period.

Conclusions

The QoL in older individuals is significantly impacted by both ADL and loneliness. In this study, the decline in ADL, coupled with advancing age and the experience of loneliness, led to a reduction in QoL among the elderly. Consequently, the subsequent recommendations are proposed:

1. For older individuals who experience loneliness, interventions that aim to strengthen social support and social contacts (e.g., visits and daily contacts) may be beneficial.
2. The development of recreational sports programs should be personalized to the needs of older adults to enhance physical activity through collaboration between health and social institutions.
3. Establishment and training on the utilization of existing infrastructure and programs for older adults need to be carried out by organizations responsible for their health and welfare.

Acknowledgments

The research proposal was accepted by the Student Research Committee, Tabriz University of Medical Sciences (Registration code: 69003). Also, the authors acknowledge the support of the "Clinical Research Development Unit, Al-Zahra Hospital," Tabriz University of Medical Sciences.

Authors contributions

Conceptualization: Mehdi Abbasian, Ehsan Sarbazi.
Investigation: Ehsan Sarbazi, Masouma Ghasemi, Mehdi Abbasian.
Project administration: Mehdi Abbasian, Ehsan Sarbazi.
Supervision: Hassan Soleimanpour.
Writing—original draft, Mehdi Abbasian, Ehsan Sarbazi, Zahra Rostami, Masouma Ghasemi.
Writing—review & editing: Ehsan Sarbazi, Mehdi Abbasian, Zahra Rostami, Hassan Soleimanpour, Masouma Ghasemi, Hosein Azizi, Mohammad Sadra Nemat, Hadi Mostafaei.

Data availability statement

All data generated or analyzed during this study are included in this published article.

Ethical approval

The Research Ethics Committee at Tabriz University of Medical Sciences reviewed and approved this study (Ethical ID: IR.TBZMED.REC.1400.1235). Additionally, oral informed consent was obtained from all individual participants included in the study.

Consent for publication

Not applicable.

Conflict of interests

The researchers of this study have no conflict of interests.

References

1. Andalib Kourayem M, Mahmoodi Nia S. The quality of life in the elderly: the role of resiliency and mindfulness. *Aging Psychol.* 2021;7(2):179-67. doi: 10.22126/jap.2021.6217.1510. [Persian].
2. World Health Organization (WHO). WHOQOL: Measuring Quality of Life. WHO; 2012. Available from: <https://www.who.int/tools/whoqol>.
3. Molzahn A, Kalfoss M, Schick-Makaroff K, Skevington SM. Comparing the importance of different aspects of quality of life to older adults across diverse cultures. *Age Ageing.* 2011;40(2):192-9. doi: 10.1093/ageing/afq156.
4. Kazemi N, Sajjadi H, Bahrami G. Quality of life in Iranian elderly. *Iran J Ageing.* 2019;13(5):518-33. doi: 10.32598/SIJA.13.Special-Issue.518. [Persian].
5. Phillips CD, Morris JN, Hawes C, Fries BE, MorV, Nennstiel M, et al. Association of the Resident Assessment Instrument (RAI) with changes in function, cognition, and psychosocial status. *J Am Geriatr Soc.* 1997;45(8):986-93. doi: 10.1111/j.1532-5415.1997.tb02971.x.
6. Chan A, Matchar DB, Tsao MA, Harding S, Chiu CT, Tay B, et al. Self-Care for Older People (SCOPE): a cluster randomized controlled trial of self-care training and health outcomes in low-income elderly in Singapore. *Contemp Clin Trials.* 2015;41:313-24. doi: 10.1016/j.cct.2015.01.001.
7. Landi F, Abbatecola AM, Provinciali M, Corsonello A, Bustacchini S, Manigrasso L, et al. Moving against frailty: does physical activity matter? *Biogerontology.* 2010;11(5):537-45. doi: 10.1007/s10522-010-9296-1.
8. Pashmdarfard M, Azad A. Assessment tools to evaluate activities of daily living (ADL) and instrumental activities of daily living (IADL) in older adults: a systematic review. *Med J Islam Repub Iran.* 2020;34:33. doi: 10.34171/mjiri.34.33.
9. Jefferson AL, Cahn-Weiner D, Boyle P, Paul RH, Moser DJ,

- Gordon N, et al. Cognitive predictors of functional decline in vascular dementia. *Int J Geriatr Psychiatry*. 2006;21(8):752-4. doi: [10.1002/gps.1556](https://doi.org/10.1002/gps.1556).
10. Medhi GK, Sarma J, Pala S, Bhattacharya H, Bora PJ, Visi V. Association between health-related quality of life (HRQOL) and activity of daily living (ADL) among elderly in an urban setting of Assam, India. *J Family Med Prim Care*. 2019;8(5):1760-4. doi: [10.4103/jfmpc.jfmpc_270_19](https://doi.org/10.4103/jfmpc.jfmpc_270_19).
 11. Christiansen L, Sanmartin Berglund J, Lindberg C, Anderberg P, Skär L. Health-related quality of life and related factors among a sample of older people with cognitive impairment. *Nurs Open*. 2019;6(3):849-59. doi: [10.1002/nop2.265](https://doi.org/10.1002/nop2.265).
 12. Habibi Sola A, Nikpour S, Seiedoshohadaei M, Haghani HA. Quality of life and status of physical functioning among elderly people in west region of Tehran: a cross-sectional survey. *Iran Journal of Nursing*. 2008;21(53):29-39. [Persian].
 13. Thakur N, Han CY. A review of assistive technologies for activities of daily living of elderly. *ArXiv [Preprint]*. June 23, 2021. Available from: <https://arxiv.org/abs/2106.12183>.
 14. Maresova P, Javanmardi E, Barakovic S, Barakovic Husic J, Tomsone S, Krejcar O, et al. Consequences of chronic diseases and other limitations associated with old age - a scoping review. *BMC Public Health*. 2019;19(1):1431. doi: [10.1186/s12889-019-7762-5](https://doi.org/10.1186/s12889-019-7762-5).
 15. Awang H, Mansor N, Nai Peng T, Nik Osman NA. Understanding ageing: fear of chronic diseases later in life. *J Int Med Res*. 2018;46(1):175-84. doi: [10.1177/0300060517710857](https://doi.org/10.1177/0300060517710857).
 16. Yousefi Z, Sameie-Sis S, Taheri-Targhi S, Araj-khodaei M, Sanaie S, Adams C, et al. Loneliness as a predictor of hypertension in older adults: the TOPS study. *Int J Aging*. 2023;1(1):e7. doi: [10.34172/ija.2023.e7](https://doi.org/10.34172/ija.2023.e7).
 17. Shearer R, Davidhizar R. It can never be the way it was: helping elderly women adjust to change and loss. *Home Healthc Nurse*. 1994;12(4):60-5. doi: [10.1097/00004045-199407000-00010](https://doi.org/10.1097/00004045-199407000-00010).
 18. Behrouz S, Saleh Moghaddam A. The Effect of Semantic Therapy on the Feeling of Loneliness of Older Women Living in Nursing Homes (Systematic Review). *Jiroft: National Conference on Aging Health*; 2021.
 19. Jylhä M. Old age and loneliness: cross-sectional and longitudinal analyses in the Tampere Longitudinal Study on Aging. *Can J Aging*. 2004;23(2):157-68. doi: [10.1353/cja.2004.0023](https://doi.org/10.1353/cja.2004.0023).
 20. Berg-Weger M, Morley J. Loneliness in old age: an unaddressed health problem. *J Nutr Health Aging*. 2020;24(3):243-5. doi: [10.1007/s12603-020-1323-6](https://doi.org/10.1007/s12603-020-1323-6).
 21. Musich S, Wang SS, Hawkins K, Yeh CS. The impact of loneliness on quality of life and patient satisfaction among older, sicker adults. *Gerontol Geriatr Med*. 2015;1:2333721415582119. doi: [10.1177/2333721415582119](https://doi.org/10.1177/2333721415582119).
 22. Hacıhasanoğlu R, Yildirim A, Karakurt P. Loneliness in elderly individuals, level of dependence in activities of daily living (ADL) and influential factors. *Arch Gerontol Geriatr*. 2012;54(1):61-6. doi: [10.1016/j.archger.2011.03.011](https://doi.org/10.1016/j.archger.2011.03.011).
 23. Miri K, Bahrami M, Vafainnya R, Gholamzadeh T. Relationship between feeling of loneliness and activities of daily living among the elderly. *Journal of Research and Health*. 2017;7(3):834-40. [Persian].
 24. Singh A, Misra N. Loneliness, depression and sociability in old age. *Ind Psychiatry J*. 2009;18(1):51-5. doi: [10.4103/0972-6748.57861](https://doi.org/10.4103/0972-6748.57861).
 25. Heravi-Karimooi M, Anoosheh M, Foroughan M, Sheykhi MT, Hajizadeh E. Understanding loneliness in the lived experiences of Iranian elders. *Scand J Caring Sci*. 2010;24(2):274-80. doi: [10.1111/j.1471-6712.2009.00717.x](https://doi.org/10.1111/j.1471-6712.2009.00717.x).
 26. Söderhamn O, Lindencrona C, Ek A. Ability for self-care among home dwelling elderly people in a health district in Sweden. *Int J Nurs Stud*. 2000;37(4):361-8. doi: [10.1016/s0020-7489\(00\)00015-8](https://doi.org/10.1016/s0020-7489(00)00015-8).
 27. Ahmadi F, Salar A, Faghihzadeh S. Assessing quality of life among elderly people in Zahedan. *Hayat*. 2004;10(22):61-7. [Persian].
 28. Montazeri A, Vahdaninia M, Mousavi SJ, Omidvari S. The Iranian version of 12-item Short Form Health Survey (SF-12): factor structure, internal consistency and construct validity. *BMC Public Health*. 2009;9:341. doi: [10.1186/1471-2458-9-341](https://doi.org/10.1186/1471-2458-9-341).
 29. Zare G, Alem J, Zare F. The validation of the measurement of the mental and physical components of SF-12 in Iranian elderly. *SHS Web Conf*. 2021;119:01002. doi: [10.1051/shsconf/202111901002](https://doi.org/10.1051/shsconf/202111901002).
 30. Ware JE, Keller SD, Kosinski M. SF-12: How to Score the SF-12 Physical and Mental Health Summary Scales. *Health Institute, New England Medical Center*; 1995.
 31. Hosseinabadi R, Foroughan M, Ghaed Amini Harouni GR, Zeidali Beiranvand R, Pournia Y. Psychometric properties of the Persian version of the 6-item De Jong Gierveld Loneliness scale in Iranian community-dwelling older persons. *Iran J Ageing*. 2020;15(3):338-49. doi: [10.32598/sija.15.3.2515.2](https://doi.org/10.32598/sija.15.3.2515.2). [Persian].
 32. Mesgar S, Amini Nasab Z, Nakhaei MH, Sharifzadeh G, Javadinia SA. Study of quality of life, depression, and daily routines in rural elders in Birjand city, Iran, in 2013. *Iran J Ageing*. 2015;10(3):142-7. [Persian].
 33. Jansson AH, Savikko NM, Pitkälä KH. Training professionals to implement a group model for alleviating loneliness among older people – 10-year follow-up study. *Educ Gerontol*. 2018;44(2-3):119-27. doi: [10.1080/03601277.2017.1420005](https://doi.org/10.1080/03601277.2017.1420005).
 34. Bond J, Bowling A, Scambler S, Victor C. Being alone in later life: loneliness, social isolation and living alone. *Revi Clin Gerontol*. 2000;10(4):407-17. doi: [10.1017/s0959259800104101](https://doi.org/10.1017/s0959259800104101).
 35. Ingrand I, Paccalin M, Liou E, Gil R, Ingrand P. Positive perception of aging is a key predictor of quality-of-life in aging people. *PLoS One*. 2018;13(10):e0204044. doi: [10.1371/journal.pone.0204044](https://doi.org/10.1371/journal.pone.0204044).
 36. Kang HW, Park M, Wallace Hernandez JP. The impact of perceived social support, loneliness, and physical activity on quality of life in South Korean older adults. *J Sport Health Sci*. 2018;7(2):237-44. doi: [10.1016/j.jshs.2016.05.003](https://doi.org/10.1016/j.jshs.2016.05.003).
 37. Macdonald B, Hülür G. Well-being and loneliness in Swiss older adults during the COVID-19 pandemic: the role of social relationships. *Gerontologist*. 2021;61(2):240-50. doi: [10.1093/geront/gnaa194](https://doi.org/10.1093/geront/gnaa194).
 38. Soleimanpour H, Sarbazi E, Davtalab Esmaeili E, Mehri A, Ghaffari Fam S, Nikbakht HA, et al. Predictors of receiving COVID-19 vaccine among adult population in Iran: an observational study. *BMC Public Health*. 2023;23(1):490. doi: [10.1186/s12889-023-15409-0](https://doi.org/10.1186/s12889-023-15409-0).
 39. Davtalab Esmaeili E, Azizi H, Sarbazi E, Khodamoradi F. The global case fatality rate due to COVID-19 in hospitalized elderly patients by sex, year, gross domestic product, and continent: a systematic review, meta-analysis, and meta-regression. *New Microbes New Infect*. 2023;51:101079. doi: [10.1016/j.nmni.2022.101079](https://doi.org/10.1016/j.nmni.2022.101079).
 40. Fakhari A, Shalchi B, Asle Rahimi V, Naghdi Sadeh R, Lak E, Najafi A, et al. Mental health literacy and COVID-19 related stress: the mediating role of healthy lifestyle in Tabriz. *Heliyon*. 2023;9(7):e18152. doi: [10.1016/j.heliyon.2023.e18152](https://doi.org/10.1016/j.heliyon.2023.e18152).
 41. Armitage R, Nellums LB. COVID-19 and the consequences of isolating the elderly. *Lancet Public Health*. 2020;5(5):e256. doi: [10.1016/s2468-2667\(20\)30061-x](https://doi.org/10.1016/s2468-2667(20)30061-x).
 42. Bonsaksen T, Thygesen H, Leung J, Ruffolo M, Schoultz M, Price D, et al. Video-based communication and its association with loneliness, mental health and quality of life among older people during the COVID-19 outbreak. *Int J Environ Res Public*

- Health. 2021;18(12):6284. doi: [10.3390/ijerph18126284](https://doi.org/10.3390/ijerph18126284).
43. Sayin Kasar K, Karaman E. Life in lockdown: social isolation, loneliness and quality of life in the elderly during the COVID-19 pandemic: a scoping review. *Geriatr Nurs*. 2021;42(5):1222-9. doi: [10.1016/j.gerinurse.2021.03.010](https://doi.org/10.1016/j.gerinurse.2021.03.010).
 44. Creese B, Khan Z, Henley W, O'Dwyer S, Corbett A, Vasconcelos Da Silva M, et al. Loneliness, physical activity, and mental health during COVID-19: a longitudinal analysis of depression and anxiety in adults over the age of 50 between 2015 and 2020. *Int Psychogeriatr*. 2021;33(5):505-14. doi: [10.1017/s1041610220004135](https://doi.org/10.1017/s1041610220004135).
 45. Minahan J, Falzarano F, Yazdani N, Siedlecki KL. The COVID-19 pandemic and psychosocial outcomes across age through the stress and coping framework. *Gerontologist*. 2021;61(2):228-39. doi: [10.1093/geront/gnaa205](https://doi.org/10.1093/geront/gnaa205).
 46. Tull MT, Edmonds KA, Scamaldo KM, Richmond JR, Rose JP, Gratz KL. Psychological outcomes associated with stay-at-home orders and the perceived impact of COVID-19 on daily life. *Psychiatry Res*. 2020;289:113098. doi: [10.1016/j.psychres.2020.113098](https://doi.org/10.1016/j.psychres.2020.113098).
 47. Bilgili N, Arpacı F. Quality of life of older adults in Turkey. *Arch Gerontol Geriatr*. 2014;59(2):415-21. doi: [10.1016/j.archger.2014.07.005](https://doi.org/10.1016/j.archger.2014.07.005).
 48. Jivraj S, Nazroo J, Vanhoutte B, Chandola T. Aging and subjective well-being in later life. *J Gerontol B Psychol Sci Soc Sci*. 2014;69(6):930-41. doi: [10.1093/geronb/gbu006](https://doi.org/10.1093/geronb/gbu006).
 49. Rababa M, Hayajneh AA, Bani-Iss W. Association of death anxiety with spiritual well-being and religious coping in older adults during the COVID-19 pandemic. *J Relig Health*. 2021;60(1):50-63. doi: [10.1007/s10943-020-01129-x](https://doi.org/10.1007/s10943-020-01129-x).
 50. Hosseini Moghaddam F, Amiri Delui M, Sadegh Moghadam L, Kameli F, Moradi M, Khajavian N, et al. Prevalence of depression and its related factors during the COVID-19 quarantine among the elderly in Iran. *Iran J Ageing*. 2021;16(1):140-51. doi: [10.32598/sija.16.1.2850.1](https://doi.org/10.32598/sija.16.1.2850.1). [Persian].
 51. Poorolajal J, Ebrahimi B, Rezapur-Shahkolai F, Doosti-Irani A, Alizadeh M, Ahmadpoor J, et al. The epidemiology of aggression and associated factors among Iranian adult population: a national survey. *J Res Health Sci*. 2020;20(4):e00499. doi: [10.34172/jrhs.2020.34](https://doi.org/10.34172/jrhs.2020.34).
 52. Arslantas D, Unsal A, Metintas S, Koc F, Arslantas A. Life quality and daily life activities of elderly people in rural areas, Eskişehir (Turkey). *Arch Gerontol Geriatr*. 2009;48(2):127-31. doi: [10.1016/j.archger.2007.11.005](https://doi.org/10.1016/j.archger.2007.11.005).
 53. Singh JA, Strand V. Spondyloarthritis is associated with poor function and physical health-related quality of life. *J Rheumatol*. 2009;36(5):1012-20. doi: [10.3899/jrheum.081015](https://doi.org/10.3899/jrheum.081015).
 54. Lee TW, Ko IS, Lee KJ. Health promotion behaviors and quality of life among community-dwelling elderly in Korea: a cross-sectional survey. *Int J Nurs Stud*. 2006;43(3):293-300. doi: [10.1016/j.ijnurstu.2005.06.009](https://doi.org/10.1016/j.ijnurstu.2005.06.009).
 55. Bidzan-Bluma I, Bidzan M, Jurek P, Bidzan L, Knietzsch J, Stueck M, et al. A Polish and German population study of quality of life, well-being, and life satisfaction in older adults during the COVID-19 pandemic. *Front Psychiatry*. 2020;11:585813. doi: [10.3389/fpsy.2020.585813](https://doi.org/10.3389/fpsy.2020.585813).
 56. Rantanen T, Eronen J, Kauppinen M, Kokko K, Sanaslahti S, Kajan N, et al. Life-space mobility and active aging as factors underlying quality of life among older people before and during COVID-19 lockdown in Finland—a longitudinal study. *J Gerontol A Biol Sci Med Sci*. 2021;76(3):e60-7. doi: [10.1093/gerona/glaa274](https://doi.org/10.1093/gerona/glaa274).
 57. Santini ZI, Jose PE, York Cornwell E, Koyanagi A, Nielsen L, Hinrichsen C, et al. Social disconnectedness, perceived isolation, and symptoms of depression and anxiety among older Americans (NSHAP): a longitudinal mediation analysis. *Lancet Public Health*. 2020;5(1):e62-70. doi: [10.1016/s2468-2667\(19\)30230-0](https://doi.org/10.1016/s2468-2667(19)30230-0).